

*Note:* Your TA probably will not cover all the problems. This is totally fine, the discussion worksheets are deliberately made long so they can serve as a resource you can use to practice, reinforce, and build upon concepts discussed in lecture, readings, and the homework.

## 1 Asymptotics and Limits

If we would like to prove asymptotic relations instead of just using them, we can use limits.

**Asymptotic Limit Rules:** If  $f(n), g(n) \geq 0$ :

- If  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} < \infty$ , then  $f(n) = \mathcal{O}(g(n))$ .
- If  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c$ , for some  $c > 0$ , then  $f(n) = \Theta(g(n))$ .
- If  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} > 0$ , then  $f(n) = \Omega(g(n))$ .

Note that these are all sufficient (and not necessary) conditions involving limits, and are not true definitions of  $\mathcal{O}$ ,  $\Theta$ , and  $\Omega$ . We highly recommend checking on your own that these statements are correct!

(a) Prove that  $n^3 = \mathcal{O}(n^4)$ .

(b) Find an  $f(n), g(n) \geq 0$  such that  $f(n) = \mathcal{O}(g(n))$ , yet  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} \neq 0$ .

(c) Prove that for any  $c > 0$ , we have  $\log n = \mathcal{O}(n^c)$ .

*Hint:* Use L'Hôpital's rule: If  $\lim_{n \rightarrow \infty} f(n) = \lim_{n \rightarrow \infty} g(n) = \infty$ , then  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{f'(n)}{g'(n)}$  (if the RHS exists)

- (d) Find an  $f(n), g(n) \geq 0$  such that  $f(n) = \mathcal{O}(g(n))$ , yet  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)}$  does not exist. In this case, you would be unable to use limits to prove  $f(n) = \mathcal{O}(g(n))$ .

*Hint: think about oscillating functions!*

## 2 Asymptotic Complexity Comparisons

- (a) Order the following functions so that for all  $i, j$ , if  $f_i$  comes before  $f_j$  in the order then  $f_i = \mathcal{O}(f_j)$ .

Do not justify your answers.

- $f_1(n) = 3^n$
- $f_2(n) = n^{\frac{1}{3}}$
- $f_3(n) = 12$
- $f_4(n) = 2^{\log_2 n}$
- $f_5(n) = \sqrt{n}$
- $f_6(n) = 2^n$
- $f_7(n) = \log_2 n$
- $f_8(n) = 2\sqrt{n}$
- $f_9(n) = n^3$

As an answer you may just write the functions as a list, e.g.  $f_8, f_9, f_1, \dots$

- (b) In each of the following, indicate whether  $f = O(g)$ ,  $f = \Omega(g)$ , or both (in which case  $f = \Theta(g)$ ). **Briefly** justify each of your answers. Recall that in terms of asymptotic growth rate, constant  $<$  logarithmic  $<$  polynomial  $<$  exponential.

	$f(n)$	$g(n)$
(i)	$\log_3 n$	$\log_4(n)$
(ii)	$n \log(n^4)$	$n^2 \log(n^3)$
(iii)	$\sqrt{n}$	$(\log n)^3$
(iv)	$n + \log n$	$n + (\log n)^2$

### 3 Recurrence Relations

Solve the following recurrence relations, assuming base cases  $T(0) = T(1) = 1$ :

(a)  $T(n) = 2 \cdot T(n/2) + O(n)$

(b)  $T(n) = T(n - 1) + n$

(c)  $T(n) = 3 \cdot T(n - 2) + 5$

(d)  $T(n) = 2 \cdot T(n/2) + O(n \log n)$

(e)  $T(n) = 3T(n^{1/3}) + O(\log n)$

(f)  $T(n) = T(n - 1) + T(n - 2)$